

RS - 5000/-

184226
29/05/2007

SEEDLING LEAF STRUCTURE OF TREE SPECIES IN
RELATION TO THEIR SUCCESSIONAL STATUS AND
CANOPY POSITION IN A SRI LANKAN RAIN FOREST.

BY

SUDUHA RUGE BANDUMALA



This thesis submitted to the University of Sri Jayewardenepura for the
award of the Degree of Master of Philosophy in Forestry on
Tropical Forest Ecology.

184226

RS - 5000/-

184226
29/05/2007

SEEDLING LEAF STRUCTURE OF TREE SPECIES IN
RELATION TO THEIR SUCCESSIONAL STATUS AND
CANOPY POSITION IN A SRI LANKAN RAIN FOREST.

BY

SUDUHAKURUGE BANDUMALA

CIR

Thesis submitted to the University of Sri Jayewardenepura for the
award of the Degree of Master of Philosophy in Forestry on
Tropical Forest Ecology.

184226


**SEEDLING LEAF STRUCTURE OF TREE SPECIES IN RELATION TO THEIR
SUCCESSIONAL STATUS AND CANOPY POSITION IN A SRI LANKA RAIN
FOREST.**

Suduhakuruge Bandumala

ABSTRACT

Many comparative studies have showed interspecific differences in physiological, anatomical and morphological traits of plants. These aspects may be combined, and interactively influence the establishment of tree seedlings under natural conditions. However, few studies have combined these aspects (physiology, morphology and growth performance) at the whole plant level. This study examined the seedling growth performance and leaf level anatomical, physiological and morphological differences in relation to species successional stage and mature tree crown position within a range of simulated shade environments. The findings of this study are important for developing silvicultural guidelines for restoration and sustainable management of tropical rain forests.

For this study, three canopy species, four subcanopy species, seven understorey species, and six pioneer species were selected. All coexist in the rain forest of southwest Sri Lanka. Seedlings were grown for two years in replicated shade houses which were designed to create shade treatments that represented a range of light quantity (photosynthetic photon flux density (PPFD) and quality (red : far red ratios) found within the Sinharaja forest. Seedling height and mortality were recorded at three months intervals. After one and half years of seedling growth, leaf photosynthesis and stomatal conductivity were measured and leaf cross sections taken for anatomical measurements. Digital graphs of leaves were taken for leaf morphology measurements. After two years of growth, seedlings were uprooted and dried at 80 °C and dry mass recorded for root, stem and leaves.



Mass ratios (Mass of plant part divided by total mass) were calculated for leaves (LMR), roots (RMR), and stem (SMR). Results showed significant differences in seedlings of climax and pioneer species in growth morphology, leaf physiology, and anatomy. Pioneer species have higher plasticity values for seedling growth (height increment, root collar diameter increment, total dry mass, stem mass ratio, root mass ratio), leaf anatomy (leaf blade thickness, upper epidermis, lower epidermis) leaf physiology (net photosynthesis, stomatal conductance) and leaf morphological characters (leaf number, specific leaf area, and stomatal density) in relation to variation in shade. This study also revealed that patterns of various seedling growth, leaf anatomical, morphological, and physiological attributes that were related to the mature tree canopy position of the seedlings. Canopy species showed higher plasticity values for height increment, leaf mass ratio, upper epidermis, net photosynthesis, leaf dry mass, leaf area and specific leaf area. Subcanopy species had higher plasticity values for root collar diameter, total dry mass, stem mass ratio, root mass ratio, palisade layer thickness and lower epidermis. Understorey species showed the lowest plasticity values for most of the measured attributes. It can be concluded that medium shade ($350 \mu\text{molm}^{-2}\text{s}^{-1}$) and light shade ($800 \mu\text{molm}^{-2}\text{s}^{-1}$) favour the optimum growth of most of the species studied..

TABLE OF CONTENTS

TABLE OF CONTENTS.....	i
LIST OF TABLES.....	iv
LIST OF FIGURES.....	vii
LIST OF PLATES.....	x
ACKNOWLEDGEMENTS.....	xv
ABSTRACT.....	xviii
CHAPTER 1: INTRODUCTION.....	1
1.1 General Introduction and Objectives.....	2
CHAPTER 2: LITERATURE REVIEW.....	9
2.1 Tropical rain forest.....	10
2.2 Plants responses to gaps.....	12
2.3 Light environment.....	15
2.4 Seedling growth performance.....	18
2.5 Leaf anatomy.....	21
2.6 Plant physiology.....	23
2.7 Seedling morphology.....	28
CHAPTER 3: STUDY DESCRIPTIONS.....	33
3.1 Study objective and hypotheses.....	34
3.2 Study site	35
3.3 Study species	36
CHAPTER 4: METHODS AND MATERIALS	40

	4.1 Experimental design.....	41
	4.2 Seedling establishment in the shelters.....	41
	4.3 Seedling growth performance experiments.....	44
	4.4 Leaf anatomy experiments.....	45
	4.5 Leaf physiology experiments.....	45
	4.6 Leaf morphology experiment.....	47
CHAPTER 5:	SEEDLING GROWTH PERFORMANCE RESULTS.....	49
	5.1 Seedling growth performance.....	50
	5.2 Seedling growth.....	55
	5.3 Whole seedling dry weight gain and allocation to roots.....	70
CHAPTER 6:	LEAF ANATOMY RESULTS.....	99
	6.0 Leaf Anatomy.....	100
	6.1 Leaf blade structure.....	100
	6.2. Stomatal density and pore length	123
CHAPTER 7:	LEAF PHYSIOLOGY RESULTS.....	136
	7.0 Seedling physiology.....	137
	7.1 Photosynthesis.....	137
	7.2 Stomatal conductance.....	147
	7.3 Nitrogen use efficiency.....	156
CHAPTER 8:	LEAF MORPHOLOGY RESULTS.....	160
	8.0 Seedling morphology.....	161
	8.1 Seedling leaf morphology.....	161
CHAPTER 9:	DISCUSSION.....	187

	9.1 Difference in seedling growth performance.....	188
	9.2 Difference in leaf anatomy.....	190
	9.3 Difference in leaf physiology.....	191
	9.4 Difference in leaf morphology.....	193
	9.5 Species differences.....	195
CHAPTER 10:	CONCLUSION.....	207
REFERENCES	210
APPENDICES 01	PHOTOGRAPHS	225
APPENDICES 02	PUBLICATIONS	267

