

SAAPCON- 2016

**5th Biennial Conference of South Asian Association of
Physiological Society of Nepal (SAAP)**

in conjunction with

**2nd Annual conference of Physiological Society of Nepal
(PSN)**

10-12 November, 2016

Kathmandu University, Dhulikhel, Nepal

Redefining Health in Nature

ABSTRACT BOOK

Organized by:



Physiological Society of Nepal (PSN)

The Opinions expressed in the abstract book do not necessarily represent the views of the Organizing Committee of SAAPCON-2016. The responsibility of opinions and for accuracy of facts presented in the abstracts rest solely on the authors. Any copying of the abstracts from this book must be done after permission from the Editorial Board (saapnepal2016@gmail.com)

Nutritional State and Neurocognitive Performance of Early Female Adolescents in Galle Municipal Area

Madhushanthi HJH¹, Wimalasekera SW², Goonewardena CSE³, Lenora RSJ⁴, Amarasekara AATD⁵

¹Allied Health Science Degree Programme, Faculty of Medicine, University of Ruhuna, Sri Lanka

²Department of Physiology, ³Department of Community Medicine, ⁴Departments of Allied Health Sciences, University of Sri Jayewardenepura, Nugegoda, Sri Lanka

Background: Nutrition is crucial for cognitive development and female adolescents are nutritionally vulnerable due to specific reasons as onset of menarche, higher requirement for growth, changes in eating pattern, life style, and risk taking behaviors.

Objective: To determine the association of nutritional status with neurocognitive functions of early female adolescents.

Methods: Cross - sectional school based study was conducted on female adolescents (11 - 14 years, n = 200). Weight, height, waist and hip circumference were measured to assess protein energy nutritional status. Tricep, bicep and subcapsular skin fold thickness (SFT) were measured to assess fat mass. Neurocognitive function was assessed with Wechsler Intelligence Tests for Children (WISC), Tests Of Nonverbal Intelligence (TONI - 3) and computerized executive function tasks. Data were analyzed by comparison of means by one way ANOVA.

Results: The nutritional categories of the sample were, normal growth (N) (29.5%), underweight (UW) (62.5 %) and overweight (OW) (8%). Mean Processing Speed Index (PSI) was significantly differed between normal 99.60(SD ±12.0) and overweight groups 78.43 SD ± 41.7) (p<0.01). Mean Working Memory Index (WMI) of normal, underweight and overweight group were 104.02 (SD ±12.0), 85.40 (SD ± 9.73) 92.75(SD ±11.24) respectively and working memory of underweight group is significantly lower than normal group (p<0.01). Abstract reasoning and executive function (inhibition and visuo spatial memory) performances were not associated with nutritional state (p>0.05).

Conclusion: Elevated BMI is negatively associated with processing speed and undernutrition is a risk factor for reduced working memory performances of female adolescents. Matured neural pathways in early delayed adolescence may contribute to the results observed in this population.