

ANTIBODY RESPONSES, HISTOPATHOLOGICAL CHANGES AND IMMUNOHISTOCHEMICAL EVIDENCE OF CHOLINERGIC NEUROIMMUNE INTERACTIONS IN LYMPHOID TISSUES IN MICE IMMUNIZED WITH LACTOCOCCUS LACTIS EXPRESSING A MALARIA PARASITE PROTEIN

Surangi Yasawardene¹, Mythreye Thayabaran² and Ranjan Ramasamy³

¹ University of Sri Jayewardenepura, Sri Lanka

² Eastern University of Sri Lanka, Sri Lanka

³ Anglia Ruskin University, United Kingdom, United Kingdom

Introduction: Gram positive food grade bacteria Lactococci have significant advantages over attenuated pathogens as vaccine delivery vehicles due to inherent safety.

Methods: Plasmodium falciparum merozoite surface antigen2(PfMSA2) was expressed in recombinant Lactococcus lactis. Recombinant L.lactis strain, delivered oro-nasally for mucosal immunisation to Balb/c mice. Non-recombinant L.lactis used as control. Serum antibody response investigated by ELISA and immunofluorescence assay. Histopathological changes in gut associated lymphoid tissue investigated. The cholinergic nerve ending and the distribution of alpha subunits of nAChRs in the spleen, Peyer's patches and mesenteric lymph nodes were studied by indirect immunohistochemistry

Results: Serum IgG anti-MSA2 responses was significantly higher in Balb/c mice, after oronasal delivery. Antibodies elicited reacted with native MSA2 in the surface of P. falciparum merozoites in an immunofluorescence assay. Enlargement of mesenteric lymph nodes and increased lymphatic infiltration of the lamina propria were noted. Spleen showed periarteriolar lymphoid aggregation in immunized mice. The $\alpha 1$ nAChRs were distributed in the capsule and red pulp of spleen. Germinal centres of Peyer's patches showed diffuse (+1) distribution of $\alpha 1$ nAChR and expressed a low immunoreactivity(IR) of anti-nAChRs.

Conclusions: Recombinant L. lactis is a suitably safe vector for subunit vaccines. Neuronal control of immune response by parasympathetic cholinergic innervation through nAChRs is suggestive.