

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/318982722>

Effect of Citrate Encapsulated and Curcumin Encapsulated Layered Double Hydroxides on in-vitro Pseudomonas aeruginosa Biofilm Growth

Conference Paper · August 2017

CITATIONS

0

READS

16

5 authors, including:



Gayan Kanchana Wijesinghe
University of Campinas

11 PUBLICATIONS 10 CITATIONS

[SEE PROFILE](#)



Ayomi Dilhari
University of Sri Jayewardenepura

14 PUBLICATIONS 14 CITATIONS

[SEE PROFILE](#)



Gd Buddhika Gayani
University of Sri Jayewardenepura

7 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



Manjula Weerasekera
University of Sri Jayewardenepura

65 PUBLICATIONS 130 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Development of graphene based devices [View project](#)



Synthesis of Magnetite Nanoparticles by Top-Down Approach from a High Purity Ore [View project](#)



ABSTRACTS



ALLIED HEALTH SCIENCES



General Sir John Kotelawala Defence University
Sri Lanka

10TH INTERNATIONAL RESEARCH CONFERENCE

3rd - 4th August 2017

'Changing Dynamics in the Global Environment: Challenges and Opportunities'

EFFECT OF LABORATORY CULTURE MEDIA, CITRATE ENCAPSULATED AND CURCUMIN ENCAPSULATED LAYERED DOUBLE HYDROXIDES ON IN-VITRO PSEUDOMONAS AERUGINOSA BIOFILM GROWTH

GK Wijesinghe¹, KAA Dilhari², GDB Gayani^{3,5}, S Kumarage³,
N Kottegoda^{3,4,5}, MM Weerasekera^{2,4#}

¹Department of Medical Laboratory Sciences, Faculty of Allied Health Sciences, General Sir John Kotelawala Defence University, Sri Lanka

²Department of Microbiology, Faculty of Medical Sciences, University of Sri Jayewardenepura, Sri Lanka

³Department of Chemistry, Faculty of Applied Sciences, University of Sri Jayewardenepura, Sri Lanka

⁴Sri Lanka Institute of Nanotechnology, Nano Science and Technology Park, Sri Lanka

⁵Advanced Material Research Center, Faculty of Applied Sciences, University of Sri Jayewardenepura, Sri Lanka

*mmwecra@sjp.ac.lk

Objective of this study was to determine the efficacy of four routine laboratory culture media on biofilm formation of *Pseudomonas aeruginosa* and the antibiofilm effect of two layered double hydroxide (LDH) nano hybrids. Influence of culture medium on *P. aeruginosa* (ATCC®27853™) adhesion as the first step of biofilm formation in the presence of four culture media (Nutrient Broth (NB), Brain Heart Infusion (BHI) broth, Luria-Bertani (LB) broth and RPMI 1640) was quantified using MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) assay after 90 minutes adhesion. *P. aeruginosa* biofilms were developed and the growth was quantified using MTT metabolic activity at 48 hour time intervals in the presence of four culture media. The effect of citrate encapsulated LDH (concentrations ranged from 1×10^{-5} g/mL to 1 g/mL) and curcumin encapsulated LDH (concentrations ranging from 1×10^{-3} g/mL to 2 g/mL) nano-hybrids on biofilms were determined using sterile 96-well microtiter plate biofilm model. The MBIC₅₀ and killing time for matured biofilms

were determined by MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay. Scanning Electron Microscopy (SEM) was performed to assess the ultra-structural changes. *P. aeruginosa* exhibited their maximum adhesion in the presence of RPMI 1640. Biofilms exhibited the maximum growth in BHI broth during 96 hour experiment period. MBIC₅₀ for curcumin-LDH was 0.1 g/mL and Citrate-LDH was 0.01 g/mL. Killing time of curcumin-LDH and Citrate-LDH were 6-12 h and less than 3h, respectively. SEM images confirmed MTT readings. The maximum planktonic and biofilm growth was achieved with BHI broth. Bacterial adhesion was enhanced in the presence of RPMI 1640. The curcumin and Citrate intercalated LDHs have a potential antibiofilm activity against *P. aeruginosa*. Further, there is a maximum antibiofilm activity against their matured biofilms within a short period of time of the treatment (3-12hours).

Keywords: Biofilms, Antibiofilm, Encapsulated layered double hydroxides